

A New Analysis of $B_6^{(1/2)}$, $B_8^{(3/2)}$, and the $\Delta I = 1/2$ Rule in the $1/N_c$ Expansion for $K \rightarrow \pi\pi$ Decays *

Gerhard O. Köhler

Institut für Physik, Universität Dortmund, D-44221 Dortmund, Germany

Abstract

We present a new calculation of long-distance contributions to the $K \rightarrow \pi\pi$ amplitudes using the $1/N_c$ expansion within the framework of chiral perturbation theory. Along these lines we compute the chiral loop corrections to the operators Q_1 , Q_2 , Q_6 , and Q_8 . In the numerical analysis we present values for $B_6^{(1/2)}$ and $B_8^{(3/2)}$. We discuss also the implications for a theoretical understanding of the $\Delta I = 1/2$ selection rule.

1 Introduction

Within the standard model the calculation of the $K \rightarrow \pi\pi$ decay amplitudes is based on the effective low-energy hamiltonian for $\Delta S = 1$ transitions [1],

$$\mathcal{H}_{eff}^{\Delta S=1} = \frac{G_F}{\sqrt{2}} \xi_u \sum_{i=1}^8 c_i(\mu) Q_i(\mu) \quad (\mu < m_c) , \quad (1)$$

$$c_i(\mu) = z_i(\mu) + \tau y_i(\mu) , \quad \tau = -\xi_t/\xi_u , \quad \xi_q = V_{qs}^* V_{qd} , \quad (2)$$

where the Wilson coefficient functions $c_i(\mu)$ of the local four-fermion operators $Q_i(\mu)$ are obtained using perturbative renormalization group techniques [2, 3]. Q_1 - Q_5 and Q_7 are composed of products of two currents, Q_6 and Q_8 are density-density operators. In view of phenomenological applications Q_1 and Q_2 , as well as, Q_6 and Q_8 are of special importance, where

$$Q_1 = 4 \bar{s}_L \gamma^\mu d_L \bar{u}_L \gamma_\mu u_L , \quad Q_2 = 4 \bar{s}_L \gamma^\mu u_L \bar{u}_L \gamma_\mu d_L , \quad (3)$$

$$Q_6 = -8 \sum_{q=u,d,s} \bar{s}_L q_R \bar{q}_R d_L , \quad Q_8 = -12 \sum_{q=u,d,s} e_q \bar{s}_L q_R \bar{q}_R d_L , \quad (4)$$

*Talk presented at the XVI Autumn School and Workshop on Fermion Masses, Mixing and CP Violation, Lisboa, Portugal, 6-15 October 1997.

with $q_{L,R} = \frac{1}{2}(1 \pm \gamma_5)q$ and $e_q = (2/3, -1/3, -1/3)$. The hadronic matrix elements of the operators, $\langle Q_i(\mu) \rangle_I \equiv \langle \pi\pi, I | Q_i(\mu) | K^0 \rangle$, are difficult to calculate but can be estimated using non-perturbative techniques generally for μ around a scale of 1 GeV. In the following analysis we will use the $1/N_c$ expansion within the framework of the chiral effective lagrangian for pseudoscalar mesons.

The calculation of chiral loop effects motivated by the $1/N_c$ expansion was introduced in Ref. [4] to investigate the $\Delta I = 1/2$ selection rule. Experimentally one finds $Re A_0 / Re A_2 \simeq 22.2$ for the ratio of the isospin amplitudes. Keeping only the dominant terms the theoretical expression for this ratio reads

$$\frac{Re A_0}{Re A_2} \simeq \frac{z_1(\mu)\langle Q_1(\mu) \rangle_0 + z_2(\mu)\langle Q_2(\mu) \rangle_0 + z_6(\mu)\langle Q_6(\mu) \rangle_0}{z_1(\mu)\langle Q_1(\mu) \rangle_2 + z_2(\mu)\langle Q_2(\mu) \rangle_2}. \quad (5)$$

It was recognized, long ago, that the vacuum saturation approximation (VSA) [5] completely fails in explaining the $\Delta I = 1/2$ rule. The authors of Ref. [4] considered chiral loop corrections to the current-current operators, Q_1 and Q_2 , and included the gluon penguin operator Q_6 at the tree level. As a net result they obtained a large enhancement of the $\Delta I = 1/2$ decay amplitude which, however, was found to be still below the experimental value ($Re A_0 = 3.33 \times 10^{-7}$ GeV). One might note that the agreement with experiment is not improved by including the NLO values for the z_i [6].

The $\Delta I = 1/2$ selection rule is governed by Q_1 and Q_2 , while the ratio ε'/ε which measures the direct CP violation in $K \rightarrow \pi\pi$ decays is dominated by the density-density operators Q_6 and Q_8 , that is to say, approximately¹

$$\varepsilon'/\varepsilon \propto y_6\langle Q_6 \rangle_0 - y_8\langle Q_8 \rangle_2 + \text{'}\pi\text{-}\eta\text{-}\eta'\text{ mixing'}. \quad (6)$$

It is of special importance to investigate whether the $1/N_c$ (chiral loop) corrections significantly affect the large cancellation between the gluon and the electroweak penguin contributions obtained at the tree level in Ref. [8].

In this talk we present a new analysis of the hadronic matrix elements of the operators Q_1 , Q_2 , Q_6 , and Q_8 in which we include the first order corrections in the twofold expansion in powers of external momenta, p , and the ratio $1/N_c$. Our main improvement concerns the matching of short- and long-distance contributions to the decay amplitude, by adopting a modified identification of virtual momenta in the chiral loops. In Section 2 we briefly review the framework of the effective low-energy approach and discuss the matching of short- and long-distance contributions. In Section 3 we list the analytic expressions for the hadronic matrix elements. Finally, in Sections 4 and 5 we present our numerical results.

We will focus here on the computation of the $K \rightarrow \pi\pi$ matrix elements. A more detailed discussion of the general method and the operator evolution at long distances is presented by T. Hambye and P. Soldan in these proceedings, and some additional details can be found in Refs. [9, 10].

¹Recent reviews on CP violation in $K \rightarrow \pi\pi$ decays can be found in Ref. [7].

2 General Framework

We start from the chiral effective lagrangian for pseudoscalar mesons which involves an expansion in momenta where terms up to $O(p^4)$ are included [11]. Keeping only terms which contribute to the $K \rightarrow \pi\pi$ matrix elements and are leading in N_c it reads

$$\begin{aligned} \mathcal{L}_{eff} = & \frac{f^2}{4} \left(\langle D_\mu U^\dagger D^\mu U \rangle + \frac{\alpha}{4N_c} \langle \ln U^\dagger - \ln U \rangle^2 + r \langle \mathcal{M} U^\dagger + U \mathcal{M}^\dagger \rangle \right) + r^2 H_2 \langle \mathcal{M}^\dagger \mathcal{M} \rangle \\ & + r L_5 \langle D_\mu U^\dagger D^\mu U (\mathcal{M}^\dagger U + U^\dagger \mathcal{M}) \rangle + r L_8 \langle \mathcal{M}^\dagger U \mathcal{M}^\dagger U + \mathcal{M} U^\dagger \mathcal{M} U^\dagger \rangle, \end{aligned} \quad (7)$$

with $D_\mu U = \partial_\mu U - i r_\mu + i U l_\mu$, $\langle A \rangle$ denoting the trace of A and $M = \text{diag}(m_u, m_d, m_s)$. f and r are free parameters related to the pion decay constant F_π and to the quark condensate, respectively, with $r = -2\langle \bar{q}q \rangle / f^2$. The complex matrix U is a non-linear representation of the pseudoscalar meson nonet:

$$U = \exp \frac{i}{f} \Pi, \quad \Pi = \pi^a \lambda_a, \quad \langle \lambda_a \lambda_b \rangle = 2\delta_{ab}. \quad (8)$$

In terms of the physical states

$$\Pi = \begin{pmatrix} \pi^0 + \frac{1}{\sqrt{3}}a\eta + \sqrt{\frac{2}{3}}b\eta' & \sqrt{2}\pi^+ & \sqrt{2}K^+ \\ \sqrt{2}\pi^- & -\pi^0 + \frac{1}{\sqrt{3}}a\eta + \sqrt{\frac{2}{3}}b\eta' & \sqrt{2}K^0 \\ \sqrt{2}K^- & \sqrt{2}\bar{K}^0 & -\frac{2}{\sqrt{3}}b\eta + \sqrt{\frac{2}{3}}a\eta' \end{pmatrix}, \quad (9)$$

with

$$a = \cos \theta - \sqrt{2} \sin \theta, \quad \sqrt{2}b = \sin \theta + \sqrt{2} \cos \theta, \quad (10)$$

and the $\eta - \eta'$ mixing angle θ for which we take $\theta = -19^\circ$ [12]. α is the instanton parameter which reflects the explicit breaking of the $U_A(1)$. The symmetry violating term must be included when treating the singlet η_0 as a dynamical degree of freedom.

The bosonic representations of the quark currents and densities are defined in terms of (functional) derivatives of the chiral action and the lagrangian, respectively:

$$\begin{aligned} \bar{q}_{iL} \gamma^\mu q_{jL} &\equiv \frac{\delta S}{\delta (l_\mu(x))_{ij}} = -i \frac{f^2}{2} (U^\dagger \partial^\mu U)_{ji} \\ &+ i r L_5 (\partial^\mu U^\dagger \mathcal{M} - \mathcal{M}^\dagger \partial^\mu U + \partial^\mu U^\dagger U \mathcal{M}^\dagger U - U^\dagger \mathcal{M} U^\dagger \partial^\mu U)_{ji}, \end{aligned} \quad (11)$$

$$\bar{q}_{iR} q_{jL} \equiv -\frac{\delta \mathcal{L}_{eff}}{\delta \mathcal{M}_{ij}} = -r \left(\frac{f^2}{4} U^\dagger + L_5 \partial_\mu U^\dagger \partial^\mu U U^\dagger + 2r L_8 U^\dagger \mathcal{M} U^\dagger + r H_2 \mathcal{M}^\dagger \right)_{ji}, \quad (12)$$

and the right-handed expressions are obtained by parity transformation. Eqs. (11) and (12) allow us to express the operators Q_i in terms of the pseudoscalar meson fields.

The $1/N_c$ corrections to the matrix elements $\langle Q_i \rangle_I$ are calculated by chiral loop diagrams. The diagrams are ultraviolet divergent and are regularized by a finite cutoff.

This procedure, which was introduced in Ref. [4], is necessary in order to restrict the chiral lagrangian to the low-energy domain. Since we truncate the effective theory to pseudoscalar mesons, the cutoff has to be taken at or, preferably, below the $\mathcal{O}(1 \text{ GeV})$.

To calculate the amplitudes we follow the lines of Ref. [4] and identify the ultraviolet cutoff of the long-distance terms with the short-distance renormalization scale μ . In carrying out this matching we pay special attention to the definition of the momenta inside the loop. This question must be addressed because the loop integrals, within the cutoff regularization, are not momentum translation invariant.

In the existing studies of the hadronic matrix elements the color singlet boson connecting the two densities or currents was integrated out from the beginning [4, 8, 13, 14]. Thus the integration variable was taken to be the momentum of the meson in the loop, and the cutoff was the upper limit of its momentum. As there is no corresponding quantity in the short-distance part, in this treatment of the integrals there is no clear matching with QCD. The ambiguity is removed, for non-factorizable diagrams, by considering the two densities to be connected to each other through the exchange of the color singlet boson, as was already discussed in Refs. [10, 15, 16, 17, 18]. A consistent matching is then obtained by assigning the same momentum to the color singlet boson at long and short distances and by identifying this momentum with the loop integration variable (see Fig. 1). The effect of this procedure is to modify the loop integrals, which introduces noticeable effects in the final results. More important it provides an unambiguous matching of the $1/N_c$ expansion in terms of mesons to the QCD expansion in terms of quarks and gluons.




Fig. 1. Matching of short- and long-distance contributions.

The approach followed here leads to an explicit classification of the diagrams into factorizable and non-factorizable: there is no way to establish a unique momentum prescription for the factorizable terms. However, factorizable loop diagrams refer to the strong sector of the theory and give corrections whose scale dependence is absorbed in the renormalization of the chiral effective lagrangian. This statement is obvious in the case of the (conserved) currents and was proved explicitly, within the cutoff formalism, in the case of the bosonized densities.² Consequently, in the factorizable sector the chiral loop corrections do not induce ultraviolet divergent terms, i.e., the only remaining ultraviolet structure of the matrix elements is contained in the overall factor $r^2 \sim 1/m_s^2$, whose scale dependence exactly cancels the factorizable evolution of the density-

²For details see Ref. [9] and the contributions of T. Hambye and P. Soldan to these proceedings.

density operators Q_6 and Q_8 at short distances. The finite terms arising from the $1/N_c$ corrections to these operators, on the other hand, will not be absorbed completely in the renormalization of the various parameters. Nevertheless, as the complete sum of the factorizable diagrams is finite, we are allowed to send the scale of the *factorizable* terms to arbitrary high values, i.e., we are allowed to calculate the remaining finite corrections in dimensional regularization, which is momentum translation invariant and yields an unambiguous result for the factorizable matrix elements. Finally, the non-factorizable loop diagrams have to be matched to the Wilson coefficients and should cancel scale dependences which arise from the (non-factorizable) short-distance expansion.

3 Hadronic Matrix Elements

Expanding the lagrangian of Eq. (7), as well as, the densities and currents of Eqs. (11) and (12) in terms of the pseudoscalar fields, the various factorizable and non-factorizable contributions to the $K \rightarrow \pi\pi$ matrix elements can be calculated from the diagrams of Figs. 2 and 3, respectively.




Fig. 2. Factorizable diagrams for the matrix elements of the operators Q_i . Crossed circles represent the currents or densities, black circles denote strong vertices. The lines represent the pseudoscalar mesons.




Fig. 3. Non-factorizable diagrams for the matrix elements of the operators Q_i .

In addition, the wave function renormalization must be included, and the values

of the various parameters in the lagrangian must be expressed in terms of physical quantities. This was done using dimensional regularization as discussed in the previous section. We verified that the scale dependence arising from the factorizable diagrams is completely canceled by the scale dependence of the various parameters in the tree level expression of the matrix elements. Finally, we obtain the following expressions:

$$i\langle\pi^+\pi^-|Q_1|K^0\rangle^F = i\langle\pi^0\pi^0|Q_2|K^0\rangle^F = 0, \quad (13)$$

$$i\langle\pi^0\pi^0|Q_1|K^0\rangle^F = -i\langle\pi^+\pi^-|Q_2|K^0\rangle^F = -X \left(1 + \frac{4L_5^r}{F_\pi^2}m_\pi^2\right), \quad (14)$$

$$i\langle\pi^0\pi^0|Q_6|K^0\rangle^F = i\langle\pi^+\pi^-|Q_6|K^0\rangle^F = -X \frac{4L_5^r}{F_\pi^2} \left(\frac{2m_K^2}{\hat{m} + m_s}\right)^2, \quad (15)$$

$$i\langle\pi^0\pi^0|Q_8|K^0\rangle^F = X \frac{2L_5^r}{F_\pi^2} \left(\frac{2m_K^2}{\hat{m} + m_s}\right)^2, \quad (16)$$

$$i\langle\pi^+\pi^-|Q_8|K^0\rangle^F = \frac{3}{4}\sqrt{2}F_\pi \left(\frac{2m_K^2}{\hat{m} + m_s}\right)^2 \left[1 + \frac{4}{3F_\pi^2}(8m_K^2 - 11m_\pi^2)L_5^r - \frac{16}{F_\pi^2}(m_K^2 - 2m_\pi^2)L_8^r\right]. \quad (17)$$

with $X = \sqrt{2}F_\pi(m_K^2 - m_\pi^2)$ and the couplings L_5^r, L_8^r defined through the relations

$$\frac{F_K}{F_\pi} = 1 + \frac{4L_5^r}{F_\pi^2}(m_K^2 - m_\pi^2), \quad (18)$$

$$\frac{m_K^2}{m_\pi^2} = \frac{\hat{m} + m_s}{2\hat{m}} \left[1 - \frac{8(m_K^2 - m_\pi^2)}{F_\pi^2}(L_5^r - 2L_8^r)\right]. \quad (19)$$

For the reason of brevity, in the analytic expressions of this section we omit scale-independent (finite) terms resulting from the loop corrections which, nevertheless, are taken into account within the numerical analysis. Note that in Eqs. (14)–(19) we used $1/F_\pi$ and $2m_K^2/(\hat{m} + m_s)$ rather than the bare parameters $1/f$ and r . Formally, the difference represents higher order effects. Nevertheless, the appearance of f or r in Eqs. (14)–(19) would induce a scale dependence which has no counterpart at the short-distance level and will be absorbed at the next order in the chiral expansion.

The non-factorizable diagrams of Fig. 3 are calculated by introducing a cutoff regulator Λ_c . Using the matching prescription of Fig. 1 we obtain

$$i\langle\pi^0\pi^0|Q_1|K^0\rangle^{NF} = 0, \quad (20)$$

$$i\langle\pi^+\pi^-|Q_1|K^0\rangle^{NF} = \frac{X}{16\pi^2 F_\pi^2} \left(-3\Lambda_c^2 + \frac{1}{4}(m_K^2 + 12m_\pi^2) \ln \Lambda_c^2\right), \quad (21)$$

$$i\langle\pi^0\pi^0|Q_2|K^0\rangle^{NF} = \frac{X}{16\pi^2 F_\pi^2} \left(\frac{9}{2}\Lambda_c^2 + \frac{3}{4}(m_K^2 - 6m_\pi^2) \ln \Lambda_c^2\right), \quad (22)$$

$$i\langle\pi^+\pi^-|Q_2|K^0\rangle^{NF} = \frac{X}{16\pi^2 F_\pi^2} \left(\frac{3}{2}\Lambda_c^2 + (m_K^2 - \frac{3}{2}m_\pi^2) \ln \Lambda_c^2\right), \quad (23)$$

$$i\langle\pi^0\pi^0|Q_6|K^0\rangle^{NF} = i\langle\pi^+\pi^-|Q_6|K^0\rangle^{NF} = \frac{X}{16\pi^2 F_\pi^2} \frac{3}{4} \left(\frac{2m_K^2}{\hat{m} + m_s}\right)^2 \ln \Lambda_c^2, \quad (24)$$

$$i\langle\pi^0\pi^0|Q_8|K^0\rangle^{NF} = \frac{X}{16\pi^2 F_\pi^2} \frac{3}{4} \left(\frac{2m_K^2}{\hat{m} + m_s}\right)^2 \ln \Lambda_c^2, \quad (25)$$

$$i\langle\pi^+\pi^-|Q_8|K^0\rangle^{NF} = -\frac{\sqrt{2}}{16\pi^2 F_\pi} \left(\frac{2m_K^2}{\hat{m} + m_s}\right)^2 \frac{\alpha}{2} \ln \Lambda_c^2. \quad (26)$$

The scale dependent terms in Eqs. (21)–(26) have to be matched to the Wilson coefficient functions. To this end we use the numerical values presented in the leading logarithm analysis of Ref. [8].

4 $B_6^{(1/2)}$ and $B_8^{(3/2)}$

To evaluate the hadronic matrix elements of Q_6 and Q_8 we pursued the following strategy. First, the non-factorizable contributions were calculated, in the isospin limit, from the diagrams of Fig. 3. In this part of the analysis the finite terms, which are systematically determined by the momentum prescription of Fig. 1, are completely taken into account. By using algebraic relations all integrals resulting from the diagrams of Fig. 3 can be reduced to three basic integrals which were calculated in the cutoff formalism. Second, the finite terms arising from the factorizable loop diagrams of Fig. 2, as well as, from the renormalization of the wave functions, the masses and the low-energy couplings were estimated using dimensional regularization, as discussed at the end of Section 2.

The values we obtain for the matrix elements of Q_6 and Q_8 are presented in Table 1, where we also specify the various contributions coming from the factorizable and the non-factorizable diagrams, respectively. In the numerical analysis we used the ratio $m_s/\hat{m} = 24.4 \pm 1.5$ [19]. In the matrix elements we extracted the factor $R^2 = [2m_K^2/(\hat{m} + m_s)]^2$, whose dependence on the (factorizable) scale is canceled exactly, for a general density-density operator, by the diagonal evolution of the Wilson coefficients.

In Table 2 we list the corresponding values for the B_i factors which quantify the deviation of the hadronic matrix elements from the VSA results:

$$B_6^{(1/2)} = |\langle Q_6 \rangle_0 / \langle Q_6 \rangle_0^{\text{VSA}}|, \quad B_8^{(3/2)} = |\langle Q_8 \rangle_2 / \langle Q_8 \rangle_2^{\text{VSA}}|. \quad (27)$$

The VSA expressions for the matrix elements were taken from Eqs. (XIX.16) and (XIX.24) of Ref. [6]. Numerically, they are $|\langle Q_6 \rangle_0| = 35.2 \cdot R^2 \text{ MeV}$ and $|\langle Q_8 \rangle_2| = 56.6 \text{ MeV} \cdot [R^2 - (0.389 \text{ GeV})^2]$. The second term in the expression for Q_8 contributes at the 2% level and has been neglected in Table 2.

We discuss next the corrections to the matrix elements $\langle Q_6 \rangle_0$ and $\langle Q_8 \rangle_2$. As already mentioned, the operator Q_6 is special because the $\mathcal{O}(p^0)$ tree level matrix element is zero due to the unitarity of the matrix U . Nevertheless the one-loop corrections to

	$\Lambda_c = 0.6 \text{ GeV}$	$\Lambda_c = 0.7 \text{ GeV}$	$\Lambda_c = 0.8 \text{ GeV}$	$\Lambda_c = 0.9 \text{ GeV}$
$i\langle Q_6 \rangle_0^{\text{tree} + \text{F loops}}$	$-68.4 - 37.0i$	$-68.4 - 37.0i$	$-68.4 - 37.0i$	$-68.4 - 37.0i$
$i\langle Q_6 \rangle_0^{\text{NF loops}}$	$29.8 + 37.0i$	$34.6 + 37.0i$	$39.0 + 37.0i$	$42.9 + 37.0i$
$ \langle Q_6 \rangle_0 ^{\text{total}}$	38.6	33.7	29.4	25.5
$i\langle Q_8 \rangle_2^{\text{tree} + \text{F loops}}$	$56.0 - 0.1i$	$56.0 - 0.1i$	$56.0 - 0.1i$	$56.0 - 0.1i$
$i\langle Q_8 \rangle_2^{\text{NF loops}}$	$-20.7 - 11.5i$	$-24.8 - 11.5i$	$-28.8 - 11.5i$	$-32.8 - 11.5i$
$ \langle Q_8 \rangle_2 ^{\text{total}}$	37.2	33.2	29.5	25.9

Table 1: Hadronic matrix elements of Q_6 and Q_8 (in units of $R^2 \cdot \text{MeV}$), shown for various values of the cutoff Λ_c .

	$\Lambda_c = 0.6 \text{ GeV}$	$\Lambda_c = 0.7 \text{ GeV}$	$\Lambda_c = 0.8 \text{ GeV}$	$\Lambda_c = 0.9 \text{ GeV}$
$B_6^{(1/2)}$	1.10	0.96	0.84	0.72
$B_8^{(3/2)}$	0.66	0.59	0.52	0.46

Table 2: B_6 and B_8 factors for various values of Λ_c .

this matrix element must be computed. These corrections are of $\mathcal{O}(p^0/N_c)$ and are non-vanishing. We have shown in Eqs. (21)–(26) that the explicit calculation of the loops yields a cutoff dependence (i.e., a non-trivial scale dependence) from the non-factorizable diagrams which has to be matched to the short-distance contribution. We note that in the twofold expansion in p^2 and p^0/N_c the contribution of the loops over the $\mathcal{O}(p^0)$ matrix element must be treated at the same level as the leading non-vanishing tree contribution proportional to L_5 . This is revealed by the large size of the non-factorizable $\mathcal{O}(p^0/N_c)$ corrections presented in Table 1. It is the sum of both, the factorizable and the non-factorizable contributions, which renders the numerical values for $\langle Q_6 \rangle_0$ close to the VSA value. The main effect of the loop corrections is to change the dependence of $\langle Q_6 \rangle_0$ on Λ_c , from a flat behaviour at the tree level to the dependence presented in Tables 1 and 2. We note that at $\Lambda_c \simeq 700 \text{ MeV}$ the value for the matrix element of Q_6 is very close to the VSA result leading to $B_6 \simeq 1$.

The Q_8 operator is not chirally suppressed, i.e., its $\mathcal{O}(p^0)$ tree level matrix element is non-zero. The one-loop corrections, even though suppressed by a factor $1/N_c$ with respect to the leading tree level, are found to be large and negative, leading to the small values for B_8 presented in Table 2. These large corrections persist in the octet limit [i.e, in the absence of the η_0 , with $a = b = 1$ and $m_\eta^2 = (4m_K^2 - m_\pi^2)/3$]. In comparison

with $\langle Q_6 \rangle_0$, the non-factorizable corrections to $\langle Q_8 \rangle_2$ are less pronounced, as expected from the power counting scheme in p^2 and $1/N_c$. However, because the factorizable $\mathcal{O}(p^2)$ and $\mathcal{O}(p^0/N_c)$ corrections are small (and negative), the non-factorizable terms produce a significant reduction of $\langle Q_8 \rangle_2$. The size of the higher order terms indicates that the leading- N_c calculation or the closely related VSA are not sufficient for the matrix elements of the Q_8 operator.

It is interesting to compare our results with those of other calculations. Refs. [13] and [14] investigated $1/N_c$ corrections to the matrix elements of Q_6 and Q_8 . This calculation considered the product of the two densities without the color singlet boson and the matching of short- and long-distance contributions was not explicit as in the present analysis. The numerical results showed also a tendency of reducing $\langle Q_8 \rangle_2$ substantially, whereas $\langle Q_6 \rangle_0$ was found to be enhanced compared to the VSA result. Calculations in lattice QCD obtain values for B_6 close to the VSA, $B_6^{(1/2)}(2 \text{ GeV}) = 1.0 \pm 0.2$ [20, 21] and $0.76(3)(5)$ [22]. Recent values reported for B_8 are $B_8^{(3/2)}(2 \text{ GeV}) = 0.81(3)(3)$ [23], $0.77(4)(4)$ [24], and $1.03(3)$ [25]. The chiral quark model [26] yields a range for B_6 which is above the VSA value, $B_6^{(1/2)}(0.8 \text{ GeV}) = 1.2 - 1.9$, and predicts a small reduction of the B_8 factor, $B_8^{(3/2)}(0.8 \text{ GeV}) = 0.91 - 0.94$. Although the scales used in the lattice calculations and the phenomenological approaches are different, the various results can be compared as the B_6 and B_8 parameters were shown in QCD to depend only very weakly on the renormalization scale for values above 1 GeV [2].

We note that our result for $B_6^{(1/2)}$ is in rough agreement with those of the various studies quoted above, whereas the value we obtain for $B_8^{(3/2)}$ lies below the values reported previously. It is desirable to investigate whether this substantial reduction of $\langle Q_8 \rangle_2$, which is due to the non-factorizable $1/N_c$ corrections to the leading term in the chiral expansion of Q_8 , will be affected by higher order corrections. This point is of great phenomenological interest because a less effective cancellation between the Q_6 and Q_8 operators, in the range obtained in the present analysis, will lead to a large value of ε'/ε in the ball park of $\sim 10^{-3}$. This can be confirmed or disproved by the forthcoming experiments at CERN (NA48), Fermilab (E832) and Frascati (KLOE).

5 $\Delta I = 1/2$ Rule

Along the lines discussed in Section 2 we computed also the hadronic matrix elements of the current-current operators Q_1 and Q_2 , including the finite terms omitted in the analytic expressions of Section 3.³ Subsequently, we carried out the matching with the short-distance coefficient functions $z_i(\mu)$. Our numerical result for the amplitude A_0 is depicted in Fig. 4. It shows an additional enhancement compared to the result of Ref. [4] which renders the amplitude in good agreement with the observed value. The new contribution arises from the Q_1 and Q_2 operators. It is due to the modified

³Note that due to current conservation no finite terms arise from the factorizable loop diagrams of Fig. 2.

matching prescription in the non-factorizable sector [except for approximately 20 % of it explained by the choice of the physical value F_π in Eqs.(21)–(26)].




Fig. 4. $Re A_0$ in units of 10^{-3} MeV for $m_s(1 \text{ GeV}) = 150 \text{ MeV}$ as a function of the matching scale Λ_c . The experimental value is represented by the dashed line.

Our result is widely stable with respect to the matching scale. The main uncertainty displayed in Fig. 4 originates from the dependence of the Wilson coefficients on Λ_{QCD} . On the other hand, the $\Delta I = 3/2$ amplitude depicted in Fig. 4 is highly unstable. The large uncertainty can be understood from the fact that it is obtained from the difference of two large amplitudes of approximately the same size [namely $A(K^0 \rightarrow \pi^+\pi^-)$ and $A(K^0 \rightarrow \pi^0\pi^0)$]. Consequently, the $\Delta I = 3/2$ amplitude is not well reproduced except that, and this is an important point, it turns out to be sufficiently suppressed whatever the particular chosen scale is between 500 MeV and 1 GeV.

In conclusion, it is certainly premature to say that the dynamical mechanism behind the $\Delta I = 1/2$ selection rule for $K \rightarrow \pi\pi$ decays is now completely understood since the $1/N_c$ expansion we use is only an approximate realization of non-perturbative QCD. In particular, vector mesons and higher resonances should be included in order to make it possible to choose a higher value for the matching scale (it is probable that the vector mesons play an important role for A_2). Nevertheless we believe that the enhancement reported here is a further important indication that the $1/N_c$ approach can account for the bulk of the $\Delta I = 1/2$ rule for $K \rightarrow \pi\pi$ decays.




Fig. 5. $Re A_2$ in units of 10^{-3} MeV for $m_s(1 \text{ GeV}) = 150 \text{ MeV}$ as a function of the matching scale Λ_c . The experimental value is represented by the dashed line.

Acknowledgements

This work has been done in collaboration with T. Hambye, E.A. Paschos, P. Soldan, and W.A. Bardeen. We wish to thank J. Bijnens and J.-M. Gérard for helpful comments. Financial support from the Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie (BMBF), 057D093P(7), Bonn, FRG, and DFG Antrag PA-10-1 is gratefully acknowledged.

References

- [1] M.K. Gaillard and B.W. Lee, Phys. Rev. Lett. **33**, (1974) 108; G. Altarelli and L. Maiani, Phys. Lett. **B52**, (1974) 351; F.J. Gilman and M.B. Wise, Phys. Rev. **D20**, (1979) 2392; B. Guberina and R.D. Peccei, Nucl. Phys. **B163**, (1980) 289.
- [2] A.J. Buras, M. Jamin, M.E. Lautenbacher, and P.H. Weisz, Nucl. Phys. **B400**, (1993) 37, 75; A.J. Buras, M. Jamin, and M.E. Lautenbacher, Nucl. Phys. **B408**, (1993) 209.
- [3] M. Ciuchini, E. Franco, G. Martinelli, and L. Reina, Phys. Lett. **B301**, (1993) 263.

- [4] W.A. Bardeen, A.J. Buras, and J.-M. Gérard, Nucl. Phys. **B293**, (1987) 787; Phys. Lett. **B192**, (1987) 138; **B211**, (1988) 343.
- [5] M.K. Gaillard and B.W. Lee, Phys. Rev. **D10**, (1974) 897.
- [6] G. Buchalla, A.J. Buras, and M.E. Lautenbacher, Rev. Mod. Phys. **68**, (1996) 1125.
- [7] A.J. Buras and R. Fleischer, in *Heavy Flavours II*, World Scientific, eds. A.J. Buras and M. Lindner (hep-ph 9704376); S. Bertolini, M. Fabbrichesi, and J.O. Eeg, report SISSA-19-98-EP (hep-ph/9802405).
- [8] G. Buchalla, A.J. Buras, and M.K. Harlander, Nucl. Phys. **B337**, (1990) 313.
- [9] T. Hambye, G.O. Köhler, E.A. Paschos, P. Soldan and W.A. Bardeen, preprint DO-TH 97/28, FERMILAB-Pub-98/051-T (hep-ph/9802300), to appear in Phys. Rev. **D**.
- [10] T. Hambye, Talk given at the 21th School of Theoretical Physics at Ustron, Poland, September 19-24 1997, Acta. Phys. Polon. **B28**, 2479-2488.
- [11] J. Gasser and H. Leutwyler, Nucl. Phys. **B250**, (1985) 465, 517, 539.
- [12] J.-M. Gérard, Mod. Phys. Lett. **A5**, (1990) 391.
- [13] J.-M. Schwarz *Diploma thesis*, Dortmund 1991; J. Heinrich and J.-M. Schwarz, *Internal Report Univ. of Dortmund* 1992-01; J. Heinrich, E.A. Paschos, J.-M. Schwarz, and Y.L. Wu, Phys. Lett. **B279**, (1992) 140.
- [14] E.A. Paschos, in *17th International Symposium on Lepton-Photon Interactions: LP95*, Proceedings, Beijing, China, 1995, edited by Zhi-Peng Zheng and He-Sheng Chen (World Scientific, Singapore, 1996).
- [15] W.A. Bardeen, talk presented at the Workshop on Hadronic Matrix Elements and Weak Decays, Ringberg Castle, Germany (April 1988), published in Nucl. Phys. Proc. Suppl. **7A**, (1989) 149.
- [16] J.P. Fatelo and J.-M. Gérard, Phys. Lett. **B347**, (1995) 136.
- [17] J. Bijnens, J.-M. Gérard, and G. Klein, Phys. Lett. **B257**, (1991) 191.
- [18] P.H. Soldan, talk presented at the Workshop on K Physics, Orsay, France (May 1996), published in the proceedings (hep-ph/9608281).
- [19] H. Leutwyler, Phys. Lett. **B378**, (1996) 313.
- [20] G. Kilcup, in *Lattice '90*, Proceedings of the International Symposium, Tallahassee, Florida, edited by V.M. Heller *et al.* [Nucl. Phys. **B (Proc. Suppl.) 20**, (1991) 417].
- [21] S. Sharpe, in *Lattice '90*, Proceedings of the International Symposium, Tallahassee, Florida, edited by V.M. Heller *et al.* [Nucl. Phys. **B (Proc. Suppl.) 20**, (1991) 429].
- [22] D. Pekurovsky and G. Kilcup, in *Lattice '97*, Proceedings of the International Symposium, Edinburgh, Scotland, edited by C. Davies *et al.*, Nucl. Phys. **B (Proc. Suppl.) 63**, (1998) 293.
- [23] T. Bhattacharya, R. Gupta, and S. Sharpe, Phys. Rev. **D55**, (1997) 4036.
- [24] G. Kilcup, R. Gupta, and S. Sharpe, Phys. Rev. **D57**, (1998) 1654.
- [25] L. Conti, A. Donini, V. Gimenez, G. Martinelli, M. Talevi, and A. Vladikas, Phys. Lett. **B421**, (1998) 273.
- [26] S. Bertolini, J.O. Eeg, M. Fabbrichesi, and E.I. Lashin, Nucl. Phys. **B514**, (1998) 93.